

## **Product datasheet for TA810174**

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TLR8 Mouse Monoclonal Antibody [Clone ID: OTI4F3]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI4F3
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 32-153 of human

TLR8(NP\_619542) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 119.6 kDa

**Gene Name:** toll like receptor 8

Database Link: NP 619542

Entrez Gene 51311 Human

Q9NR97

Synonyms: CD288


Protein Families: Druggable Genome, Transmembrane

**Protein Pathways:** Toll-like receptor signaling pathway






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TLR8 (Cat# [RC221986], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TLR8 antibody (Cat# TA810174)(1:2000). Positive lysates [LY408461] (100ug) and [LC408461] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human spleen tissue within the normal limits using anti-TLR8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810174) (1:2000)





Immunohistochemical staining of paraffinembedded Carcinoma of Human spleen tissue within the normal limits using anti-TLR8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810174}) (1:2000).