

Product datasheet for TA810174M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TLR8 Mouse Monoclonal Antibody [Clone ID: OTI4F3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4F3
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 32-153 of human

TLR8(NP_619542) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 119.6 kDa

Gene Name: toll like receptor 8

Database Link: NP 619542

Entrez Gene 51311 Human

Q9NR97

Synonyms: CD288


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Toll-like receptor signaling pathway






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TLR8 (Cat# [RC221986], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TLR8 antibody (Cat# [TA810174])(1:2000). Positive lysates [LY408461] (100ug) and [LC408461] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human spleen tissue within the normal limits using anti-TLR8 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Carcinoma of Human spleen tissue within the normal limits using anti-TLR8 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.