

Product datasheet for TA810249M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GTP cyclohydrolase 1 (GCH1) Mouse Monoclonal Antibody [Clone ID: OTI8C6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8C6
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:500 Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GCH1 (NP_001019195) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27.7 kDa

Gene Name: GTP cyclohydrolase 1

Database Link: NP 001019195

Entrez Gene 14528 MouseEntrez Gene 29244 RatEntrez Gene 2643 Human

P30793

Synonyms: DYT5; DYT5a; DYT14; GCH; GTP-CH-1; GTPCH1; HPABH4B


Protein Families: Druggable Genome

Protein Pathways: Folate biosynthesis, Metabolic pathways






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GCH1 ([RC224968], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GCH1 (1:2000). Positive lysates [LY422611] (100ug) and [LC422611] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-GCH1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-GCH1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.