

Product datasheet for TA810269S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MLH1 Mouse Monoclonal Antibody [Clone ID: OTI5A7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5A7

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 299-527 of human

MLH1 (NP_000240) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 84.4 kDa

Gene Name: mutL homolog 1

Database Link: NP 000240

Entrez Gene 17350 MouseEntrez Gene 81685 RatEntrez Gene 4292 Human

P40692

Background: This gene was identified as a locus frequently mutated in hereditary nonpolyposis colon

cancer (HNPCC). It is a human homolog of the E. coli DNA mismatch repair gene mutL, consistent with the characteristic alterations in microsatellite sequences (RER+phenotype) found in HNPCC. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described, but their full-length natures

have not been determined. [provided by RefSeq, Nov 2009]




MLH1 Mouse Monoclonal Antibody [Clone ID: OTI5A7] - TA810269S

Synonyms: COCA2; FCC2; hMLH1; HNPCC; HNPCC2; MMRCS1

Protein Families: Druggable Genome

Protein Pathways: Colorectal cancer, Endometrial cancer, Mismatch repair, Pathways in cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MLH1 ([RC201607], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MLH1 (1:500). Positive lysates [LY400096] (100ug) and [LC400096] (20ug) can be purchased separately from OriGene.