

Product datasheet for TA810308BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PGBD3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3H1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3H1

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 313-593 of human

PGBD3 (NP_736609) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: piggyBac transposable element derived 3

Database Link: NP 736609

Entrez Gene 267004 Human

Q8N328






Background:

This gene is a member of a small family of genes derived from piggyBac transposable elements. The encoded protein contains a zinc-ribbon domain characteristic of transposon-derived proteins and may function as a regulator of transcription. Alternative splicing occurs between a splice site from exon 5 of the adjacent upstream gene 'excision repair cross-complementation group 6' (ERCC6, GeneID: 2074) and the 3' splice site upstream of the open reading frame (ORF) of this gene, which activates the alternative polyadenylation site downstream of the piggyback-derived-3 ORF. The resulting transcripts encode a fusion protein that shares sequence with the product of each individual gene. Pseudogenes for this gene are defined on chromosomes 4, 5 and 12. [provided by RefSeq, Mar 2016]

Synonyms: FLJ90201

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PGBD3 ([RC208977], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PGBD3 (1:2000). Positive lysates [LY406870] (100ug) and [LC406870] (20ug) can be purchased separately from OriGene.