

Product datasheet for TA810334M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BAX Mouse Monoclonal Antibody [Clone ID: OTI2A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A1

Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-172 of human BAX

(NP 620116) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21 kDa

Gene Name: BCL2 associated X protein

Database Link: NP 620116

Entrez Gene 12028 MouseEntrez Gene 24887 RatEntrez Gene 581 Human

Q07812

Synonyms: BCL2L4

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Amyotrophic lateral sclerosis (ALS), Apoptosis, Colorectal cancer, Huntington's disease,


Neurotrophin signaling pathway, p53 signaling pathway, Pathways in cancer, Prion diseases





Product images:







HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY BAX (Cat# [RC204369], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BAX (1:2000) (Cat# [TA810334]). Positive lysates [LY408504] (100ug) and [LC408504] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 6 different cell lines by using anti-BAX monoclonal antibody (1:500).






Equivalent amounts of cell lysates (10 ug per lane) of wild-type Hela cells (WT, Cat# LC810HELA) and BAX-Knockout Hela cells (KO, Cat# [LC810020]) were separated by SDS-PAGE and immunoblotted with anti-BAX monoclonal antibody [TA810334]. Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control (1:500).



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-BAX mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-BAX mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.