

Product datasheet for TA810357BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CISD2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4F8
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CISD2 (NP_001008389) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: CDGSH iron sulfur domain 2

Database Link: NP 001008389

Entrez Gene 67006 MouseEntrez Gene 295457 RatEntrez Gene 493856 Human

Q8N5K1


Synonyms: ERIS; Miner1; NAF-1; WFS2; ZCD2

Protein Families: Transmembrane






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CISD2 ([RC207131], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CISD2 (1:2000).




Western blot analysis of extracts (35ug) from 8 different cell lines and rat brain tissue lysate by using anti-CISD2 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-CISD2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810357]) (1:150)



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CISD2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810357]) (1:150)