

## **Product datasheet for TA810475S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MED19 Mouse Monoclonal Antibody [Clone ID: OTI4C3]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4C3

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human MED19 (NP\_703151) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 20.3 kDa

**Gene Name:** mediator complex subunit 19

Database Link: NP 703151

Entrez Gene 311165 RatEntrez Gene 381379 MouseEntrez Gene 219541 Human

A0JLT2

**Background:** MED19 is a component of the Mediator complex, which is a coactivator for DNA-binding

factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed

12584197]). [supplied by OMIM, Oct 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC037223.1, EL733643.1 [ECO:0000332] RNAseq introns :: single sample supports all introns

ERS025082, ERS025095 [ECO:0000348] ##Evidence-Data-END##






**Synonyms:** DT2P1G7; LCMR1; MED19AS

**Protein Families:** Druggable Genome

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MED19 ([RC206806], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MED19 (1:2000). Positive lysates [LY407036] (100ug) and [LC407036] (20ug) can be purchased separately from OriGene.