

# **Product datasheet for TA810565M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **RUNX3 Mouse Monoclonal Antibody [Clone ID: OTI11D5]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI11D5
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:500 Human, Mouse, Rat

**Host:** Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human RUNX3 (NP\_001026850) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 45.7 kDa

**Gene Name:** runt related transcription factor 3

Database Link: NP 001026850

Entrez Gene 12399 MouseEntrez Gene 156726 RatEntrez Gene 864 Human

Q13761


Synonyms: AML2; CBFA3; PEBP2aC

Protein Families: Transcription Factors






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RUNX3 ([RC203597], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RUNX3 (1:2000). Positive lysates [LY400418] (100ug) and [LC400418] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-RUNX3 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-RUNX3 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.