

# **Product datasheet for TA810598**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **CPT1B Mouse Monoclonal Antibody [Clone ID: OTI6B8]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6B8
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:500 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 187-440 of human

CPT1B (NP 004368) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 87.6 kDa

**Gene Name:** carnitine palmitoyltransferase 1B

Database Link: NP 004368

Entrez Gene 12895 MouseEntrez Gene 25756 RatEntrez Gene 1375 Human

092523

Synonyms: CPT1-M; CPT1M; CPTI-M; M-CPT1; MCCPT1; MCPT1


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Adipocytokine signaling pathway, Fatty acid metabolism, PPAR signaling pathway






## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CPT1B ([RC217297], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CPT1B (1:2000). Positive lysates [LY418031] (100ug) and [LC418031] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-CPT1B mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810598) (1:500)