

## **Product datasheet for TA810618M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **BAG5 Mouse Monoclonal Antibody [Clone ID: OTI3C10]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3C10

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BAG5 (NP\_004864) produced in 293T.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** BCL2 associated athanogene 5

Database Link: NP 004864

Entrez Gene 70369 MouseEntrez Gene 366734 RatEntrez Gene 9529 Human

Q9UL15

**Background:** The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an

anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. Three transcript variants encoding two different isoforms have been

found for this gene. [provided by RefSeq, Jul 2008]

Synonyms: BAG-5







**Protein Families:** 

Druggable Genome

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BAG5 ([RC208518], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BAG5 (1:2000). Positive lysates [LY417691] (100ug) and [LC417691] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 7 different cell lines by using anti-BAG5 monoclonal antibody (1:500).