

### Product datasheet for TA810791AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PGAM4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A1
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombiant protein of human PGAM4 (NP\_001025062) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 28.6 kDa

**Gene Name:** phosphoglycerate mutase family member 4

Database Link: NP 001025062

Entrez Gene 441531 Human

**Q8N0Y7** 


Synonyms: dJ1000K24.1; PGAM-B; PGAM1; PGAM3

**Protein Pathways:** Glycolysis / Gluconeogenesis, Metabolic pathways






# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PGAM4 ([RC213480], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PGAM4 (1:2000). Positive lysates [LY422267] (100ug) and [LC422267] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 2 different cell lines (293T and MCF7) by using anti-PGAM4 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Human adult heart tissue within the normal limits using anti-PGAM4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810791]) (1:500)