

Product datasheet for **TA810807BM**

PLAC8 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2E10]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2E10
Applications:	IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:500
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PLAC8 (NP_057703) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	12.3 kDa
Gene Name:	placenta specific 8
Database Link:	NP_057703 Entrez Gene 51316 Human Q9NZF1
Synonyms:	C15; DGIC; onzin; PNAS-144




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PLAC8 (Cat# [RC212588], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PLAC8 (Cat# [TA810807])(1:2000). Positive lysates [LY413872] (100ug) and [LC413872] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from human liver tissue lysate by using anti-PLAC8 monoclonal antibody (1:500).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-PLAC8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810807]) (1:500)



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PLAC8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810807]) (1:500)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-PLAC8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810807]) (1:500)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-PLAC8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810807]) (1:500)