

Product datasheet for TA810901BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HNRPH1 (HNRNPH1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1H3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1H3
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:500

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombiant protein of human HNRNPH1 (NP_005511) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49 kDa

Gene Name: heterogeneous nuclear ribonucleoprotein H1 (H)

Database Link: NP 005511

Entrez Gene 59013 MouseEntrez Gene 3187 Human


P31943

Synonyms: hnRNPH; HNRPH; HNRPH1






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HNRNPH1 ([RC201834], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HNRNPH1 (1:2000). Positive lysates [LY417247] (100ug) and [LC417247] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 8 different cell lines by using anti-HNRNPH1 monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)







Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)




Immunohistochemical staining of paraffinembedded Human thyroid tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810901]) (1:500)



Immunohistochemical staining of paraffinembedded Human gastric tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA (1:500)



Immunohistochemical staining of paraffinembedded Human spleen tissue within the normal limits using anti-HNRNPH1 mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA (1:500)