

# **Product datasheet for TA810972S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **HOXD8 Mouse Monoclonal Antibody [Clone ID: OTI6D10]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6D10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 69-290 of human

HOXD8 (NP\_062458) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 31.7 kDa

Gene Name: homeobox D8

Database Link: NP 062458

Entrez Gene 3234 Human

P13378






Background:

This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conserved family of transcription factors that play an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, located on different chromosomes, consisting of 9 to 11 genes arranged in tandem. This gene is one of several homeobox HOXD genes located in a cluster on chromosome 2. Deletions that remove the entire HOXD gene cluster or the 5' end of this cluster have been associated with severe limb and genital abnormalities. In addition to effects during embryogenesis, this particular gene may also play a role in adult urogenital tract function. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Dec 2010]

Synonyms: HOX4; HOX4E; HOX5.4

**Protein Families:** ES Cell Differentiation/IPS, Transcription Factors

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HOXD8 ([RC205062], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HOXD8. Positive lysates [LY412699] (100ug) and [LC412699] (20ug) can be purchased separately from OriGene.