

# Product datasheet for TA810973AM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IMPDH2 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IMPDH2 (NP\_000875) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 55.6 kDa

**Gene Name:** IMP (inosine 5'-monophosphate) dehydrogenase 2

Database Link: NP 000875

Entrez Gene 23918 MouseEntrez Gene 301005 RatEntrez Gene 3615 Human

P12268

**Background:** This gene encodes the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis.

It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. The encoded protein catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. This gene is up-regulated in some neoplasms, suggesting it

may play a role in malignant transformation. [provided by RefSeq, Jul 2008]






Synonyms: IMPD2; IMPDH-II

**Protein Families:** Druggable Genome

**Protein Pathways:** Drug metabolism - other enzymes, Metabolic pathways, Purine metabolism

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IMPDH2 (Cat# [RC202977], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IMPDH2 antibody (Cat# [TA810973]). Positive lysates [LY424469] (100ug) and [LC424469] (20ug) can be purchased separately from OriGene.