

Product datasheet for TA811050AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RNF14 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C9
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-220 of human RNF14

(NP_899647) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 53.7 kDa

Gene Name: ring finger protein 14

Database Link: NP 899647

Entrez Gene 56736 MouseEntrez Gene 619577 RatEntrez Gene 9604 Human

Q9UBS8


Synonyms: ARA54; HFB30; HRIHFB2038; TRIAD2

Protein Families: Druggable Genome, Transcription Factors






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RNF14 (Cat# [RC223362], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RNF14 (Cat# [TA811050])(1:2000). Positive lysates [LY405271] (100ug) and [LC405271] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 4 different cell lines by using anti-RNF14 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-RNF14 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811050]) (1:500)



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-RNF14 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811050]) (1:500)