

Product datasheet for TA811057M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nck beta (NCK2) Mouse Monoclonal Antibody [Clone ID: OTI4H7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4H7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NCK2 (NP_003572) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42.7 kDa

Gene Name: NCK adaptor protein 2

Database Link: NP 003572

Entrez Gene 17974 MouseEntrez Gene 316369 RatEntrez Gene 8440 Human

<u>O43639</u>

Background: This gene encodes a member of the NCK family of adaptor proteins. The protein contains

three SH3 domains and one SH2 domain. The protein has no known catalytic function but has been shown to bind and recruit various proteins involved in the regulation of receptor protein tyrosine kinases. It is through these regulatory activities that this protein is believed to be involved in cytoskeletal reorganization. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]




Synonyms: GRB4; NCKbeta

Protein Families: Druggable Genome

Protein Pathways: Axon guidance, ErbB signaling pathway, Pathogenic Escherichia coli infection, T cell receptor

signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NCK2 (Cat# [RC202557], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NCK2(Cat# [TA811057]). Positive lysates [LY401190] (100ug) and [LC401190] (20ug) can be purchased separately from OriGene.