

Product datasheet for **TA811105BM**


ABCF2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8F4]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI8F4
Applications:	IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:500
Reactivity:	Human, Mouse
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-235 of human ABCF2 (NP_009120) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	71.1 kDa
Gene Name:	ATP binding cassette subfamily F member 2
Database Link:	NP_009120 Entrez Gene 27407 Mouse Entrez Gene 10061 Human Q9UG63
Synonyms:	ABC28; EST133090; HUSSY-18; HUSSY18
Protein Families:	Druggable Genome




[View online »](#)

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ABCF2 (Cat# [RC200323], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ABCF2 (Cat# [TA811105])(1:2000). Positive lysates [LY402100] (100ug) and [LC402100] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 3 different cell lines by using anti-ABCF2 monoclonal antibody (1:500).




Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and ABCF2-Knockout 293T cells (KO, Cat# [LC811907]) were separated by SDS-PAGE and immunoblotted with anti-ABCF2 monoclonal antibody [TA811105], (1:500). Then the blotted membrane was stripped and reprobed with anti-beta-actin antibody ([TA811000]) as a loading control.



Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-ABCF2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811105]) (1:500)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-ABCF2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811105]) (1:500)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-ABCF2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811105]) (1:500)