

## **Product datasheet for TA811232AM**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RNF4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI9F10]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI9F10

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Rat, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-190 of human RNF4

(NP\_002929) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** ring finger protein 4

Database Link: NP 001171938

Entrez Gene 19822 MouseEntrez Gene 29274 RatEntrez Gene 6047 Human

P78317

**Background:** The protein encoded by this gene contains a RING finger motif and acts as a transcription

regulator. This protein has been shown to interact with, and inhibit the activity of, TRPS1, a

transcription suppressor of GATA-mediated transcription. Transcription repressor

ZNF278/PATZ is found to interact with this protein, and thus reduce the enhancement of androgen receptor-dependent transcription mediated by this protein. Studies of the mouse and rat counterparts suggested a role of this protein in spermatogenesis. A pseudogene of

this gene is found on chromosome 1. [provided by RefSeq, Jul 2010]







Synonyms: RES4-26; SLX5; SNURF

**Protein Families:** Transcription Factors

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RNF4 ([RC207273], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RNF4 (1:2000). Positive lysates [LY432659] (100ug) and [LC432659] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 5 different cell lines by using anti-RNF4 monoclonal antibody (1:500).