

Product datasheet for TA811407M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VAX1 Mouse Monoclonal Antibody [Clone ID: OTI1E6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E6
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human VAX1 (NP_954582) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 20.9 kDa

Gene Name: ventral anterior homeobox 1

Database Link: NP 954582

Entrez Gene 22326 MouseEntrez Gene 64571 RatEntrez Gene 11023 Human

Q5SQQ9


Synonyms: MCOPS11

Protein Families: Transcription Factors






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VAX1 ([RC210985], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VAX1 (1:2000). Positive lysates [LY404744] (100ug) and [LC404744] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-VAX1 mouse monoclonal antibody. This figure shows negative staining. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-VAX1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.