

Product datasheet for TA811427AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FAM54A (MTFR2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A7
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FAM54A (NP_001092756) produced in

E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.2 kDa

Gene Name: mitochondrial fission regulator 2

Database Link: NP 001092756

Entrez Gene 113115 Human


Q6P444

Synonyms: DUFD1; FAM54A






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY FAM54A (Cat# [RC213401], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FAM54A (Cat# [TA811427])(1:500). Positive lysates [LY426058] (100ug) and [LC426058] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-FAM54A mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811427]) (1:150)





Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-FAM54A mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811427]) (1:150)



Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-FAM54A mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811427]) (1:150)