

Product datasheet for TA811464

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

NEMF Mouse Monoclonal Antibody [Clone ID: OTI7E10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7E10

Applications: IHC

Recommended Dilution: IHC 1:250

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 869-1076 of human

NEMF (NP_004704) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 122.8 kDa

Gene Name: nuclear export mediator factor

Database Link: NP 004704

Entrez Gene 66244 MouseEntrez Gene 100322884 RatEntrez Gene 9147 Human


<u>O60524</u>

Synonyms: IDDSAPN; NY-CO-1; SDCCAG1






Product images:




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-NEMF mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-NEMF mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-NEMF mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-NEMF mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.