

Product datasheet for TA811481BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Peroxiredoxin 5 (PRDX5) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4B12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4B12
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PRDX5 (NP_036226) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21.8 kDa

Gene Name: peroxiredoxin 5

Database Link: NP 036226

Entrez Gene 54683 MouseEntrez Gene 113898 RatEntrez Gene 25824 Human

P30044


Synonyms: ACR1; AOEB166; B166; HEL-S-55; PLP; PMP20; PRDX6; prx-V; PRXV; SBBI10

Protein Families: Druggable Genome






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PRDX5 (Cat# [RC210709], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PRDX5 (Cat# [TA811481])(1:2000). Positive lysates [LY402144] (100ug) and [LC402144] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 6 different cell lines by using anti-PRDX5 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-PRDX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811481]) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-PRDX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811481]) (1:150)