

Product datasheet for TA811539S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD14 Mouse Monoclonal Antibody [Clone ID: OTI6D7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6D7

Applications: FC, IHC, WB

Recommended Dilution: WB 1:500, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 20-87 of human CD14

(NP_000582) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 40.08 kDa

Gene Name: CD14 molecule

Database Link: NP 000582

Entrez Gene 929 Human

P08571

Background: The protein encoded by this gene is a surface antigen that is preferentially expressed on

monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript

variants encoding the same protein. [provided by RefSeq, Mar 2010]

Synonyms: CD14 antigen; CD14 molecule; monocyte differentiation antigen CD14




Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,

Transmembrane


Protein Pathways: Hematopoietic cell lineage, MAPK signaling pathway, Pathogenic Escherichia coli infection,

Regulation of actin cytoskeleton, Toll-like receptor signaling pathway

Product images:




Western blot analysis of extracts (35ug) from PBMC by using anti-CD14 monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CD14 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811539]) (1:150)





HEK293T cells transfected with either [RC203819] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CD14 antibody ([TA811539]), and then analyzed by flow cytometry (1:100).