

# **Product datasheet for TA811544S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **FOLR3 Mouse Monoclonal Antibody [Clone ID: OTI7E10]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7E10
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human FOLR3 (NP\_000795) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** folate receptor 3 (gamma)

Database Link: NP 000795

Entrez Gene 2352 Human

P41439

Background: This gene encodes a member of the folate receptor (FOLR) family of proteins, which have a

high affinity for folic acid and for several reduced folic acid derivatives, and mediate delivery of 5-methyltetrahydrofolate to the interior of cells. Expression of this gene may be elevated in

ovarian and primary peritoneal carcinoma. This gene is present in a gene cluster on chromosome 11. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Dec 2015]

**Synonyms:** FR-G; FR-gamma; gamma-hFR







**Protein Families:** 

Druggable Genome, Secreted Protein

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FOLR3 ([RC212963], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FOLR3 (1:2000).



Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-FOLR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811544]) (1:2000)