

## Product datasheet for **TA811641AM**

### **RACGAP1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4H8]**

#### **Product data:**


<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI4H8
<b>Applications:</b>	IHC, WB
<b>Recommended Dilution:</b>	WB 1:2000, IHC 1:500
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Human recombinant protein fragment corresponding to amino acids 123-238 of human RACGAP1 (NP_001119575) produced in E.coli.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.5 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Conjugation:</b>	Biotin
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Gene Name:</b>	Rac GTPase activating protein 1
<b>Database Link:</b>	<a href="#">NP_001119575</a> <a href="#">Entrez Gene 26934 Mouse</a> <a href="#">Entrez Gene 315298 Rat</a> <a href="#">Entrez Gene 29127 Human</a> <a href="#">Q9H0H5</a>
<b>Background:</b>	This gene encodes a GTPase-activating protein (GAP) that is a component of the centralspindlin complex. This protein binds activated forms of Rho GTPases and stimulates GTP hydrolysis, which results in negative regulation of Rho-mediated signals. This protein plays a regulatory role in cytokinesis, cell growth, and differentiation. Alternatively spliced transcript variants have been found for this gene. There is a pseudogene for this gene on chromosome 12. [provided by RefSeq, Feb 2016]
<b>Synonyms:</b>	CYK4; HsCYK-4; ID-GAP; MgcRacGAP




[View online »](#)

Protein Families: Druggable Genome


### Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RACGAP1 ([RC226045], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RACGAP1 (1:2000). Positive lysates [LY426645] (100ug) and [LC426645] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Human spleen tissue within the normal limits using anti-RACGAP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811641]) (1:500)



Immunohistochemical staining of paraffin-embedded Human gastric stromal tumor tissue using anti-RACGAP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811641]) (1:500)