

### Product datasheet for TA811677AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **EEF1D Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6B3]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6B3
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human EEF1D (NP\_115754) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** eukaryotic translation elongation factor 1 delta

Database Link: NP 115754

Entrez Gene 1936 Human

P29692

**Background:** This gene encodes a subunit of the elongation factor-1 complex, which is responsible for the


enzymatic delivery of aminoacyl tRNAs to the ribosome. This subunit, delta, functions as guanine nucleotide exchange factor. It is reported that following HIV-1 infection, this subunit interacts with HIV-1 Tat. This interaction results in repression of translation of host cell proteins and enhanced translation of viral proteins. Several alternatively spliced transcript variants encoding multiple isoforms have been found for this gene. Related pseudogenes have been defined on chromosomes 1, 6, 7, 9, 11, 13, 17, 19. [provided by RefSeq, Aug 2010]


**Synonyms:** EF-1D; EF1D; FP1047






# **Product images:**









Western blot analysis of extracts (35ug) from 2 different cell lines by using anti-EEF1D monoclonal antibody (1:500).






Immunohistochemical staining of paraffinembedded Human adult brain tissue within the normal limits using anti-EEF1D mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811677]) (1:2000)



Immunohistochemical staining of paraffinembedded Human embryonic cerebellum within the normal limits using anti-EEF1D mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811677]) (1:2000)



Immunohistochemical staining of paraffinembedded Human testicle tissue within the normal limits using anti-EEF1D mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811677]) (1:2000)