

Product datasheet for TA811680S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pancreatic Lipase (PNLIP) Mouse Monoclonal Antibody [Clone ID: OTI9A9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9A9
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:1000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 355-465 of human

PNLIP (NP_000927) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: pancreatic lipase

Database Link: NP 000927

Entrez Gene 5406 Human

P16233

Background: This gene is a member of the lipase gene family. It encodes a carboxyl esterase that

hydrolyzes insoluble, emulsified triglycerides, and is essential for the efficient digestion of dietary fats. This gene is expressed specifically in the pancreas. [provided by RefSeq, Jul 2008]

Synonyms: PL; PNLIPD; PTL


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Glycerolipid metabolism, Metabolic pathways






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PNLIP ([RC204735], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PNLIP (1:2000). Positive lysates [LY424451] (100ug) and [LC424451] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-PNLIP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811680]) (1:1000)