

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA811700

## KPNB1 Mouse Monoclonal Antibody [Clone ID: OTI10D3]

## Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI10D3
Applications:	WB
Recommended Dilution:	WB 1:500~2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-200 of human KPNB1 (NP_002256) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	97 kDa
Gene Name:	karyopherin subunit beta 1
Database Link:	<u>NP_002256</u> <u>Entrez Gene 16211 MouseEntrez Gene 24917 RatEntrez Gene 3837 Human</u> <u>Q14974</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** KPNB1 Mouse Monoclonal Antibody [Clone ID: OTI10D3] – TA811700

Background:Nucleocytoplasmic transport, a signal- and energy-dependent process, takes place through<br/>nuclear pore complexes embedded in the nuclear envelope. The import of proteins<br/>containing a nuclear localization signal (NLS) requires the NLS import receptor, a heterodimer<br/>of importin alpha and beta subunits also known as karyopherins. Importin alpha binds the<br/>NLS-containing cargo in the cytoplasm and importin beta docks the complex at the<br/>cytoplasmic side of the nuclear pore complex. In the presence of nucleoside triphosphates<br/>and the small GTP binding protein Ran, the complex moves into the nuclear pore complex<br/>and the importin subunits dissociate. Importin alpha enters the nucleoplasm with its<br/>passenger protein and importin beta remains at the pore. Interactions between importin beta<br/>and the FG repeats of nucleoporins are essential in translocation through the pore complex.<br/>The protein encoded by this gene is a member of the importin beta family. Two transcript<br/>variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb<br/>2013]


Synonyms:IMB1; Impnb; IPO1; IPOB; NTF97Protein Families:Druggable Genome, Stem cell - Pluripotency

#### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KPNB1 ([RC200659], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KPNB1 (1:2000). Positive lysates [LY419417] (100ug) and [LC419417] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Western blot analysis of extracts (35ug) from 2 different cell lines by using anti-KPNB1 monoclonal antibody (1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US