

## Product datasheet for TA811713BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### IRF2BP1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI6C9]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6C9
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:250

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 457-584 of human

IRF2BP1 (NP\_056464) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 61.5 kDa

**Gene Name:** interferon regulatory factor 2 binding protein 1

Database Link: NP 056464

Entrez Gene 272359 MouseEntrez Gene 308404 RatEntrez Gene 26145 Human

Q8IU81


Synonyms: DKFZp434M154

**Protein Families:** Druggable Genome






# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IRF2BP1 ([RC207151], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IRF2. Positive lysates [LY414410] (100ug) and [LC414410] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from HepG2 cell line human liver tissue lysate by using anti-IRF2BP1 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-IRF2BP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811713]) (1:250)