

ESEARCH

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA811714

KPNB1 Mouse Monoclonal Antibody [Clone ID: OTI9D6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI9D6
Applications:	WB
Recommended Dilution:	WB 1:500~2000
Reactivity:	Human, Rat, Mouse
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-200 of human KPNB1 (NP_002256) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	97 kDa
Gene Name:	karyopherin subunit beta 1
Database Link:	<u>NP_002256</u> <u>Entrez Gene 16211 MouseEntrez Gene 24917 RatEntrez Gene 3837 Human</u> <u>Q14974</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE KPNB1 Mouse Monoclonal Antibody [Clone ID: OTI9D6] – TA811714

Background:Nucleocytoplasmic transport, a signal- and energy-dependent process, takes place through
nuclear pore complexes embedded in the nuclear envelope. The import of proteins
containing a nuclear localization signal (NLS) requires the NLS import receptor, a heterodimer
of importin alpha and beta subunits also known as karyopherins. Importin alpha binds the
NLS-containing cargo in the cytoplasm and importin beta docks the complex at the
cytoplasmic side of the nuclear pore complex. In the presence of nucleoside triphosphates
and the small GTP binding protein Ran, the complex moves into the nuclear pore complex
and the importin subunits dissociate. Importin alpha enters the nucleoplasm with its
passenger protein and importin beta remains at the pore. Interactions between importin beta
and the FG repeats of nucleoporins are essential in translocation through the pore complex.
The protein encoded by this gene is a member of the importin beta family. Two transcript
variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb
2013]


Synonyms:IMB1; Impnb; IPO1; IPOB; NTF97Protein Families:Druggable Genome, Stem cell - Pluripotency

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KPNB1 ([RC200659], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KPNB1 (1:2000). Positive lysates [LY419417] (100ug) and [LC419417] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Western blot analysis of extracts (35ug) from 4 different cell lines by using anti-KPNB1 monoclonal antibody (1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US