

## **Product datasheet for TA811720AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Dermcidin (DCD) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3F5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F5
Applications: WB

December and ad Dilutions - WD 1.50

Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 20-110 of human DCD

(NP\_444513) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 9.2 kDa

Gene Name: dermcidin

Database Link: NP 444513

Entrez Gene 117159 Human

P81605



# Dermcidin (DCD) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3F5] – TA811720AM


Background:

This antimicrobial gene encodes a secreted protein that is subsequently processed into mature peptides of distinct biological activities. The C-terminal peptide is constitutively expressed in sweat and has antibacterial and antifungal activities. The N-terminal peptide, also known as diffusible survival evasion peptide, promotes neural cell survival under conditions of severe oxidative stress. A glycosylated form of the N-terminal peptide may be associated with cachexia (muscle wasting) in cancer patients. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2014]

Synonyms: AIDD; DCD-1; DSEP; HCAP; PIF

**Protein Families:** Secreted Protein

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DCD (Cat# [RC209352], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DCD (Cat# [TA811720])(1:500). Positive lysates [LY403291] (100ug) and [LC403291] (20ug) can be purchased separately from OriGene.