

## **Product datasheet for TA811770S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **METTL3 Mouse Monoclonal Antibody [Clone ID: OTI1B7]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1B7

**Applications:** WB

Recommended Dilution: WB 1:500~2000

**Reactivity:** Human, Rat, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-284 of human

METTL3 (NP\_062826) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 64.3 kDa

**Gene Name:** methyltransferase like 3

Database Link: NP 062826

Entrez Gene 56335 MouseEntrez Gene 361035 RatEntrez Gene 56339 Human

Q86U44

Background: This gene encodes the 70 kDa subunit of MT-A which is part of N6-adenosine-

methyltransferase. This enzyme is involved in the posttranscriptional methylation of internal adenosine residues in eukaryotic mRNAs, forming N6-methyladenosine. [provided by RefSeq,


Jul 20081

Synonyms: IME4; M6A; MT-A70; Spo8






### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY METTL3 (Cat# [RC200869], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-METTL3 (Cat# [TA811770])(1:2000). Positive lysates [LY412720] (100ug) and [LC412720] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 6 different cell lines and 3 different tissue lysates by using anti-METTL3 monoclonal antibody (1:500).