

Product datasheet for TA811872BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ACSL5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1D4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1D4

Applications: WB

Reactivity: WB 1:500
Host: WB 0:500

WB 1:500

Human

Mouse

Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 104-620 of human

ACSL5 (NP_976313) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 75.8 kDa

Gene Name: acyl-CoA synthetase long-chain family member 5

Database Link: NP 976313

Entrez Gene 51703 Human

O9ULC5





Background:

The protein encoded by this gene is an isozyme of the long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme is highly expressed in uterus and spleen, and in trace amounts in normal brain, but has markedly increased levels in malignant gliomas. This gene functions in mediating fatty acid-induced glioma cell growth. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]


Synonyms: ACS2; ACS5; FACL5

Protein Families: Transmembrane

Protein Pathways: Adipocytokine signaling pathway, Fatty acid metabolism, Metabolic pathways, PPAR signaling

pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACSL5 ([RC213380], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACSL5. Positive lysates [LY404321] (100ug) and [LC404321] (20ug) can be purchased separately from OriGene.