

# **Product datasheet for TA811874S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **RFC3 Mouse Monoclonal Antibody [Clone ID: OTI3F6]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F6
Applications: IHC, WB

**Reactivity:** WB 1:500, IHC 1:2000 Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human RFC3 (NP\_002906) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 40.4 kDa

**Gene Name:** replication factor C subunit 3

Database Link: NP 002906

Entrez Gene 69263 MouseEntrez Gene 288414 RatEntrez Gene 5983 Human

P40938

**Background:** The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase

epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and

replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kDa. This gene encodes the 38 kDa subunit. This subunit is essential for the interaction between the 140 kDa subunit and the core complex that consists of the 36, 37, and 40 kDa subunits. Alternatively spliced transcript variants

encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]






Synonyms: RFC38


**Protein Families:** Stem cell - Pluripotency

**Protein Pathways:** DNA replication, Mismatch repair, Nucleotide excision repair

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RFC3 (Cat# [RC201655], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RFC3(Cat# [TA811874]). Positive lysates [LY419018] (100ug) and [LC419018] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-RFC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811874]) (1:2000)





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-RFC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA811874]) (1:2000)