

### Product datasheet for TA811907BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# G protein alpha inhibitor 1 (GNAI1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2D1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D1

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-354 of human GNAI1

(NP 002060) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 40.2 kDa

**Gene Name:** G protein subunit alpha i1

Database Link: NP 002060

Entrez Gene 14677 MouseEntrez Gene 25686 RatEntrez Gene 2770 Human

P63096





# G protein alpha inhibitor 1 (GNAI1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2D1] – TA811907BM

**Background:** Guanine nucleotide binding proteins are heterotrimeric signal-transducing molecules

consisting of alpha, beta, and gamma subunits. The alpha subunit binds guanine nucleotide, can hydrolyze GTP, and can interact with other proteins. The protein encoded by this gene represents the alpha subunit of an inhibitory complex. The encoded protein is part of a complex that responds to beta-adrenergic signals by inhibiting adenylate cyclase. Two transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jan 2012]

Synonyms: Gi


**Protein Families:** Druggable Genome

**Protein Pathways:** Axon guidance, Chemokine signaling pathway, Gap junction, Leukocyte transendothelial

migration, Long-term depression, Melanogenesis, Progesterone-mediated oocyte maturation,

Tight junction

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY GNAI1 (Cat# [RC205289], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GNAI1 (Cat# [TA811907]). Positive lysates [LY419561] (100ug) and [LC419561] (20ug) can be purchased separately from OriGene.