

Product datasheet for TA811910M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Interferon alpha 2 (IFNA2) Mouse Monoclonal Antibody [Clone ID: OTI12C7]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: OTI12C7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 24-188 of human IFNA2

(NP 000596) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

lgG1

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21.4 kDa

Gene Name: interferon, alpha 2

Database Link: NP 000596

Entrez Gene 3440 Human

P01563

Background: This gene is a member of the alpha interferon gene cluster on chromosome 9. The encoded

protein is a cytokine produced in response to viral infection. Use of the recombinant form of this protein has been shown to be effective in reducing the symptoms and duration of the

common cold. [provided by RefSeq, Jun 2011]

Synonyms: IFN-alphaA; IFNA; IFNA2B; INFA2






Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Antigen processing and presentation, Autoimmune thyroid disease, Cytokine-cytokine

receptor interaction, Cytosolic DNA-sensing pathway, Jak-STAT signaling pathway, Natural killer cell mediated cytotoxicity, Regulation of autophagy, RIG-I-like receptor signaling

pathway, Toll-like receptor signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IFNA2 (Cat# [RC221091], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFNA2(Cat# [TA811910]). Positive lysates [LY400201] (100ug) and [LC400201] (20ug) can be purchased separately from OriGene.