

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA811921BM

# PSMD5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8B1]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI8B1
Applications:	WB
Recommended Dilution:	WB 1:500~2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 2-140 of human PSMD5 (NP_005038) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	56 kDa
Gene Name:	proteasome 26S subunit, non-ATPase 5
Database Link:	<u>NP_005038</u> <u>Entrez Gene 66998 MouseEntrez Gene 296651 RatEntrez Gene 5711 Human</u> <u>Q16401</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### SMD5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8B1] – TA811921BM

Background:The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure<br/>composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4<br/>rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are<br/>composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6<br/>ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase<br/>subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration<br/>and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This<br/>gene encodes a non-ATPase subunit of the 19S regulator base that functions as a chaperone<br/>protein during 26S proteasome assembly. [provided by RefSeq, Jul 2012]

#### Synonyms:

### **Product images:**



A431


170 -130 -100 -70 -

40 -


35 — 25 — 15 — 10 — Jurkat

S5B

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PSMD5 (Cat# [RC204913], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PSMD5(Cat# [TA811921]). Positive lysates [LY401563] (100ug) and [LC401563] (20ug) can be purchased separately from OriGene.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and PSMD5-Knockout HeLa cells (KO, Cat# [LC832826]) were separated by SDS-PAGE and immunoblotted with anti-PSMD5 monoclonal antibody [TA811921] (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US