

## **Product datasheet for TA812012AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **CREM Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4H9]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4H9

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human CREM (NP\_874388) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 13.3 kDa

**Gene Name:** cAMP responsive element modulator

Database Link: NP 874388

Entrez Gene 12916 MouseEntrez Gene 25620 RatEntrez Gene 1390 Human

Q03060






### Background:

This gene encodes a bZIP transcription factor that binds to the cAMP responsive element found in many viral and cellular promoters. It is an important component of cAMP-mediated signal transduction during the spermatogenetic cycle, as well as other complex processes. Alternative promoter and translation initiation site usage allows this gene to exert spatial and temporal specificity to cAMP responsiveness. Multiple alternatively spliced transcript variants encoding several different isoforms have been found for this gene, with some of them functioning as activators and some as repressors of transcription. [provided by RefSeq, Jul 2008]

Synonyms: CREM-2; hCREM-2; ICER

**Protein Families:** Druggable Genome, Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CREM ([RC217044], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CREM. Positive lysates [LY403643] (100ug) and [LC403643] (20ug) can be purchased separately from OriGene.