

Product datasheet for TA812019S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Alpha Fodrin (SPTAN1) Mouse Monoclonal Antibody [Clone ID: OTI9C10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9C10
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-147 of human

SPTAN1 (NP_001123910) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: spectrin alpha, non-erythrocytic 1

Database Link: NP 001123910

Entrez Gene 20740 MouseEntrez Gene 64159 RatEntrez Gene 6709 Human

Q13813





Background:


Spectrins are a family of filamentous cytoskeletal proteins that function as essential scaffold proteins that stabilize the plasma membrane and organize intracellular organelles. Spectrins are composed of alpha and beta dimers that associate to form tetramers linked in a head-to-head arrangement. This gene encodes an alpha spectrin that is specifically expressed in nonerythrocytic cells. The encoded protein has been implicated in other cellular functions including DNA repair and cell cycle regulation. Mutations in this gene are the cause of early infantile epileptic encephalopathy-5. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2010]

Synonyms: DEE5; EIEE5; NEAS; SPTA2


Protein Families: Druggable Genome

Protein Pathways: Tight junction

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY SPTAN1 (Cat# [RC226485], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SPTAN1 (Cat# [TA812019])(1:2000).




Western blot analysis of extracts (35ug) from 3 cell line by using anti-SPTAN1 monoclonal antibody (1:500).






Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)




Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)



Immunohistochemical staining of paraffinembedded Human gastric tissue within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)





Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-SPTAN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812019]) (1:2000)