

### Product datasheet for TA812059BM

lgG1

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **EPO Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D8]**

#### **Product data:**

**Applications:** 

Isotype:

**Product Type:** Primary Antibodies

Clone Name: OTI5D8

Recommended Dilution:WB 1:150Reactivity:HumanHost:Mouse

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 28-193 of human EPO

(NP\_000790) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 21.31 kDa

Gene Name: erythropoietin

Database Link: NP 000790

Entrez Gene 2056 Human

P01588

**Background:** This gene is a member of the EPO/TPO family and encodes a secreted, glycosylated cytokine

composed of four alpha helical bundles. The protein is found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. This protein also has neuroprotective activity against a variety of potential brain injuries and

antiapoptotic functions in several tissue types. [provided by RefSeq, Jul 2008]




**Synonyms:** EP; MVCD2

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling

pathway

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EPO (Cat# [RC210775], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EPO (Cat# [TA812059]). Positive lysates [LY400277] (100ug) and [LC400277] (20ug) can be purchased separately from OriGene.