

Product datasheet for TA812060BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Somatostatin Receptor 5 (SSTR5) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1E4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E4

Applications: WB

Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse
Isotype: IgG2c

Clonality: Monoclonal

Immunogen: Synthetic peptide (the amino acid sequence is considered to be commercially sensitive)

within Human SSTR5(NP_001044). The exact sequence is proprietary.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: somatostatin receptor 5

Database Link: NP 001044

Entrez Gene 6755 Human

P35346





Somatostatin Receptor 5 (SSTR5) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1E4] – TA812060BM

Background:


Somatostatin and its related peptide cortistatin exert multiple biological actions on normal and tumoral tissue targets by interacting with somatostatin receptors (SSTRs). The protein encoded by this gene is one of the SSTRs, which is a multi-pass membrane protein and belongs to the G-protein coupled receptor 1 family. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase, and different regions of this receptor molecule are required for the activation of different signaling pathways. A mutation in this gene results in somatostatin analog resistance. Alternatively spliced transcript variants have been identified in this gene. [provided by RefSeq, Feb 2010]

Synonyms: SS-5-R

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Equivalent amounts of lysates (5 ug per lane) of SSTR1, 2, 3, 4 and 5 peptide (from lane 1 to 5) were separated by SDS-PAGE and immunoblotted with anti-SSTR5 antibody (1:500).