

## **Product datasheet for TA812116**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **HSPA1L Mouse Monoclonal Antibody [Clone ID: OTI3B10]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3B10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 400-641 of human

HSPA1L (NP\_005518) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 70.2 kDa

**Gene Name:** heat shock protein family A (Hsp70) member 1 like

Database Link: NP 005518

Entrez Gene 15482 MouseEntrez Gene 24963 RatEntrez Gene 3305 Human

P34931

**Background:** This gene encodes a 70kDa heat shock protein. In conjunction with other heat shock proteins,

this protein stabilizes existing proteins against aggregation and mediates the folding of newly

translated proteins in the cytosol and in organelles. The gene is located in the major

histocompatibility complex class III region, in a cluster with two closely related genes which also encode isoforms of the 70kDa heat shock protein. [provided by RefSeg, Jul 2008]






Synonyms: HSP70-1L; HSP70-HOM; HSP70T; hum70t

**Protein Pathways:** Antigen processing and presentation, Endocytosis, MAPK signaling pathway, Spliceosome

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HSPA1L ([RC205351], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSPA1. Positive lysates [LY417244] (100ug) and [LC417244] (20ug) can be purchased separately from OriGene.