

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA812187AM

# Bone marrow stromal cell antigen 1 (BST1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI9B2]

### **Product data:**

Product Type:	Primary Antibodies		
Clone Name:	OTI9B2		
Applications:	WB		
Recommended Dilution:	WB 1:2000		
Reactivity:	Human		
Host:	Mouse		
lsotype:	lgG2a		
Clonality:	Monoclonal		
Immunogen:	Human recombinant protein fragment corresponding to amino acids 29-318 of human BST1 (NP_004325) produced in E.coli.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Concentration:	0.5 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	Biotin		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	35.5 kDa		
Gene Name:	bone marrow stromal cell antigen 1		
Database Link:	<u>NP_004325</u> <u>Entrez Gene 683 Human</u> <u>Q10588</u>		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Bone marrow stromal cell antigen 1 (BST1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI9B2] – TA812187AM
Background:	Bone marrow stromal cell antigen-1 is a stromal cell line-derived glycosylphosphatidylinositol- anchored molecule that facilitates pre-B-cell growth. The deduced amino acid sequence exhibits 33% similarity with CD38. BST1 expression is enhanced in bone marrow stromal cell lines derived from patients with rheumatoid arthritis. The polyclonal B-cell abnormalities in rheumatoid arthritis may be, at least in part, attributed to BST1 overexpression in the stromal cell population. [provided by RefSeq, Jul 2008]
Synonyms:	CD157
Protein Families	: Transmembrane
Protein Pathwa	<b>ys:</b> Calcium signaling pathway, Metabolic pathways, Nicotinate and nicotinamide metabolism

## **Product images:**

170	—	
130	_	
100	_	19
70	—	
55		
40		
35	—	
25	-	
15	_	
10	—	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BST1 ([RC204151], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BST1. Positive lysates [LY418052] (100ug) and [LC418052] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US