

## Product datasheet for TA812187M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Bone marrow stromal cell antigen 1 (BST1) Mouse Monoclonal Antibody [Clone ID: OTI9B2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI9B2
Applications: WB

December de di Dilindiano - NAD 4-20

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 29-318 of human BST1

(NP\_004325) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 35.5 kDa

**Gene Name:** bone marrow stromal cell antigen 1

Database Link: NP 004325

Entrez Gene 683 Human

Q10588

Bone marrow stromal cell antigen-1 is a stromal cell line-derived glycosylphosphatidylinositol-

anchored molecule that facilitates pre-B-cell growth. The deduced amino acid sequence exhibits 33% similarity with CD38. BST1 expression is enhanced in bone marrow stromal cell lines derived from patients with rheumatoid arthritis. The polyclonal B-cell abnormalities in rheumatoid arthritis may be, at least in part, attributed to BST1 overexpression in the stromal

cell population. [provided by RefSeq, Jul 2008]






# Bone marrow stromal cell antigen 1 (BST1) Mouse Monoclonal Antibody [Clone ID: OTI9B2] – TA812187M

Synonyms: CD157

**Protein Families:** Transmembrane

**Protein Pathways:** Calcium signaling pathway, Metabolic pathways, Nicotinate and nicotinamide metabolism

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BST1 ([RC204151], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BST1. Positive lysates [LY418052] (100ug) and [LC418052] (20ug) can be purchased separately from OriGene.