

# **Product datasheet for TA812355M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **TNFRSF4 Mouse Monoclonal Antibody [Clone ID: OTI9G3]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI9G3
Applications: FC, WB

Recommended Dilution: WB 1:1000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TNFRSF4 (NP\_003318) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 29.34 kDa

**Gene Name:** tumor necrosis factor receptor superfamily member 4

Database Link: NP 003318

Entrez Gene 7293 Human

P43489

**Background:** The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of

apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The

knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul 2008]






Synonyms: ACT35; CD134; IMD16; OX40; TXGP1L


**Protein Families:** Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNFRSF4 ([RC211253], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF4 (1:1000).



Flow cytometric analysis of living 293T cells transfected with TNFRSF4 overexpression plasmid ([RC211253]), Red)/empty vector ([PS100001], Blue) using anti-TNFRSF4 antibody ([TA812355]). Cells incubated with a non-specific antibody (Green) were used as isotype control.