

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA812357M

## **TNFRSF4 Mouse Monoclonal Antibody [Clone ID: OTI1B10]**

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI1B10
Applications:	FC, WB
Recommended Dilution:	WB 1:1000, FLOW 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TNFRSF4 (NP_003318) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.34 kDa
Gene Name:	tumor necrosis factor receptor superfamily member 4
Database Link:	<u>NP_003318</u> <u>Entrez Gene 7293 Human</u> <u>P43489</u>
Background:	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul 2008]



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### Service TNFRSF4 Mouse Monoclonal Antibody [Clone ID: OTI1B10] – TA812357M

Synonyms:


Protein Families: Transmembrane

Protein Pathways:


Cytokine-cytokine receptor interaction

ACT35; CD134; IMD16; OX40; TXGP1L

#### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNFRSF4 ([RC211253], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF4 (1:1000).



Flow cytometric analysis of living 293T cells transfected with TNFRSF4 overexpression plasmid ([RC211253]), Red)/empty vector ([PS100001], Blue) using anti-TNFRSF4 antibody ([TA812357]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-TNFRSF4 antibody ([TA812357]) (1:100).

Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-TNFRSF4 antibody ([TA812357]) (Right). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US