

### **Product datasheet for TA812382**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## c Maf (MAF) Mouse Monoclonal Antibody [Clone ID: OTI4H10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4H10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human MAF (NP\_001026974) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** MAF bZIP transcription factor

Database Link: NP 001026974

Entrez Gene 17132 MouseEntrez Gene 54267 RatEntrez Gene 4094 Human

075444

**Background:** The protein encoded by this gene is a DNA-binding, leucine zipper-containing transcription

factor that acts as a homodimer or as a heterodimer. Depending on the binding site and binding partner, the encoded protein can be a transcriptional activator or repressor. This protein plays a role in the regulation of several cellular processes, including embryonic lens fiber cell development, increased T-cell susceptibility to apoptosis, and chondrocyte terminal differentiation. Defects in this gene are a cause of juvenile-onset pulverulent cataract as well as congenital cerulean cataract 4 (CCA4). Two transcript variants encoding different isoforms

have been found for this gene. [provided by RefSeq, Jan 2010]






**Synonyms:** AYGRP; c-MAF; CCA4; CTRCT21

**Protein Families:** Druggable Genome, Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MAF (Cat# [RC216495], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAF (Cat# TA812382)(1:2000).