

Product datasheet for TA812384BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DIP13B (APPL2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1H8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1H8

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human APPL2 (NP_060641) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 74.3 kDa

Gene Name: adaptor protein, phosphotyrosine interacting with PH domain and leucine zipper 2

Database Link: NP 060641

Entrez Gene 55198 Human

Q8NEU8







Background:

The protein encoded by this gene is one of two effectors of the small GTPase RAB5A/Rab5, which are involved in a signal transduction pathway. Both effectors contain an N-terminal Bin/Amphiphysin/Rvs (BAR) domain, a central pleckstrin homology (PH) domain, and a C-terminal phosphotyrosine binding (PTB) domain, and they bind the Rab5 through the BAR domain. They are associated with endosomal membranes and can be translocated to the nucleus in response to the EGF stimulus. They interact with the NuRD/MeCP1 complex (nucleosome remodeling and deacetylase /methyl-CpG-binding protein 1 complex) and are required for efficient cell proliferation. A chromosomal aberration t(12;22)(q24.1;q13.3) involving this gene and the PSAP2 gene results in 22q13.3 deletion syndrome, also known as Phelan-McDermid syndrome. [provided by RefSeq, Oct 2011]

Synonyms: DIP13B

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY APPL2 (Cat# [RC207506], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-APPL2(Cat# [TA812384]). Positive lysates [LY413268] (100ug) and [LC413268] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from cell lines and/or tissue lysates by using anti-APPL2 monoclonal antibody (1:500).